

ICOS ligand Antibody

Mouse Monoclonal Antibody [Clone ICOSL/9016]

Catalog No	Format	Size
23308-MSM13-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
23308-MSM13-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
23308-MSM13-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details	
Clone	ICOSL/9016
Immunogen	Recombinant human ICOSLG protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	33.35kDa
Cellular Localization	Cell membrane
Species Reactivity	Human
Positive Control	Expressed on peripheral blood B-cells and monocytes, as well as on monocyte-derived dendritic cells (at protein level)

*Optimal dilution for a specific application should be determined.

Product Images for ICOS ligand Antibody

Specificity & Comments

Ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion (PubMed:11007762, PubMed:11023515, PubMed:30498080). Also induces B-cell proliferation and differentiation into plasma cells. Could play an important role in mediating local tissue responses to inflammatory conditions, as well as in modulating the secondary immune response by co-stimulating memory T-cell function (By similarity). In endothelial cells, required for proper neutrophil transmigration in response to chemoattractants, such as CXCL8/IL8 or N-formyl-methionyl peptides (fMLP) (PubMed:30498080).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.