

5-hydroxytryptamine receptor 1A Antibody

Mouse Monoclonal Antibody [Clone HTR1A/15323]

Catalog No	Format	Size
3350-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3350-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3350-MSM3-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	HTR1A/15323
Immunogen	Recombinant fragment (around aa 218-345) of human HTR1A protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1
Mol. Weight of Antigen	46.11kDa
Cellular Localization	Cell membrane, Cell projection, Dendrite
Species Reactivity	Human
Positive Control	Detected in lymph nodes, thymus and spleen

*Optimal dilution for a specific application should be determined.

Product Images for 5-hydroxytryptamine receptor 1A Antibody

Specificity & Comments

G-protein coupled receptor for 5-hydroxytryptamine (serotonin) (PubMed:22957663, PubMed:3138543, PubMed:33762731, PubMed:37935376, PubMed:37935377, PubMed:8138923, PubMed:8393041). Also functions as a receptor for various drugs and psychoactive substances (PubMed:22957663, PubMed:3138543, PubMed:33762731, PubMed:38552625, PubMed:8138923, PubMed:8393041). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylyate cyclase (PubMed:22957663, PubMed:3138543, PubMed:33762731, PubMed:8138923, PubMed:8393041). HTR1A is coupled to G(i)/G(o) G alpha proteins and mediates inhibitory neurotransmission: signaling inhibits adenylyate cyclase activity and activates a phosphatidylinositol-calcium second messenger system that regulates the release of Ca(2+) ions from intracellular stores (PubMed:33762731, PubMed:35610220). Beta-arrestin family members regulate signaling by mediating both receptor desensitization and resensitization processes (PubMed:18476671, PubMed:20363322, PubMed:20945968). Plays a role in the regulation of 5-hydroxytryptamine release and in the regulation of dopamine and 5-hydroxytryptamine metabolism (PubMed:18476671, PubMed:20363322, PubMed:20945968). Plays a role in the regulation of dopamine and 5-hydroxytryptamine levels in the brain, and thereby affects neural activity, mood and behavior (PubMed:18476671, PubMed:20363322, PubMed:20945968). Plays a role in the response to anxiogenic stimuli (PubMed:18476671, PubMed:20363322, PubMed:20945968).

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.