Interleukin-1beta (IL-1beta) Antibody

Mouse Monoclonal Antibody [Clone IL1B/15413]

Catalog No	Format	Size
3553-MSM19-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3553-MSM19-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3553-MSM19-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	IL1B/15413
Immunogen	Recombinant fragment (around aa 117-269) of human IL1B protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1
Mol. Weight of Antigen	30.75kDa
Cellular Localization	Cytoplasm, Cytosol, Extracellular exosome, Lysosome, Secreted
Species Reactivity	Human
Positive Control	Expressed in activated monocytes/macrophages (at protein level)

**Optimal dilution for a specific application should be determined.*

Product Images for Interleukin-1beta (IL-1beta) Antibody

Specificity & Comments

Potent pro-inflammatory cytokine (PubMed:10653850, PubMed:12794819, PubMed:28331908, PubMed:3920526). Initially discovered as the major endogenous pyrogen, induces prostaglandin synthesis, neutrophil influx and activation, T-cell activation and cytokine production, B-cell activation and antibody production, and fibroblast proliferation and collagen production (PubMed:3920526). Promotes Th17 differentiation of T-cells. Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells (PubMed:10653850). Plays a role in angiogenesis by inducing VEGF production synergistically with TNF and IL6 (PubMed:12794819). Involved in transduction of inflammation downstream of pyroptosis: its mature form is specifically released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33377178, PubMed:33883744). Acts as a sensor of S.pyogenes infection in skin: cleaved and activated by pyogenes SpeB protease, leading to an inflammatory response that prevents bacterial growth during invasive skin infection (PubMed:28331908).

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.