Recombinant Interleukin-13 Antibody

Rabbit Monoclonal Antibody [Clone IL13/17235R]

Catalog No	Format	Size
3596-RBM16-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3596-RBM16-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3596-RBM16-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	IL13/17235R
Immunogen	Recombinant human IL13 protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	15.79kDa
Cellular Localization	Secreted
Species Reactivity	Human

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant Interleukin-13 Antibody



SDS-PAGE under non-reducing (NR) and reducing (R) conditions. The gel was stained with Coomassie Blue. The purity of the protein is 96%.

SDS-PAGE Analysis of Purified Recombinant Interleukin-13 Rabbit Monoclonal Antibody (IL13/17235R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

Cytokine that plays important roles in allergic inflammation and immune response to parasite infection (PubMed:8096327, PubMed:8097324). Synergizes with IL2 in regulating interferon-gamma synthesis (PubMed:8096327). Stimulates B-cell proliferation, and activation of eosinophils, basophils, and mast cells (PubMed:7903680, PubMed:8759755). Plays an important role in controlling IL33 activity by modulating the production of transmembrane and soluble forms of interleukin-1 receptor-like 1/IL1RL1 (By similarity). Displays the capacity to antagonize Th1-driven proinflammatory immune response and downregulates synthesis of many proinflammatory cytokines including IL1, IL6, IL10, IL12 and TNF-alpha through a mechanism that partially involves suppression of NF-kappa-B (By similarity). Also functions on nonhematopoietic cells, including endothelial cells where it induces vascular cell adhesion protein 1/VCAM1, which is important in the recruitment of eosinophils (PubMed:8639787). Exerts its biological effects through its receptors which comprises the IL4R chain and the IL13RA1 chain, to activate JAK1 and TYK2, leading to the activation of STAT6 (PubMed:9013879). Aside from IL13RA1, another receptor IL13RA2 acts as a high affinity decoy for IL13 and mediates internalization and depletion of extracellular IL13 (PubMed:21622864).

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.