

## Interleukin-18 Antibody

Mouse Monoclonal Antibody [Clone IL18/4629]

Catalog No	Format	Size
3606-MSM9-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3606-MSM9-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3606-MSM9-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note

### Product Details

<b>Clone</b>	IL18/4629
<b>Immunogen</b>	Recombinant human IL18 protein
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2 / Kappa
<b>Mol. Weight of Antigen</b>	22.33kDa
<b>Cellular Localization</b>	Cytoplasm, Cytosol, Secreted
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Expressed in ovarian carcinoma but undetectable in normal ovarian epithelial cells

\*Optimal dilution for a specific application should be determined.

### Product Images for Interleukin-18 Antibody

#### Specificity & Comments

Pro-inflammatory cytokine primarily involved in epithelial barrier repair, polarized T-helper 1 (Th1) cell and natural killer (NK) cell immune responses (PubMed:10653850). Upon binding to IL18R1 and IL18RAP, forms a signaling ternary complex which activates NF-kappa-B, triggering synthesis of inflammatory mediators (PubMed:14528293, PubMed:25500532, PubMed:37993714). Synergizes with IL12/interleukin-12 to induce IFNG synthesis from T-helper 1 (Th1) cells and natural killer (NK) cells (PubMed:10653850). Involved in transduction of inflammation downstream of pyroptosis: its mature form is specifically released in the extracellular milieu by passing through the gasdermin-D (GSDMD) pore (PubMed:33883744).

#### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

#### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

#### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.