

Inhibin, alpha (INHA) (Gonadal Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone INHA/9352]

Catalog No	Format	Size
3623-MSM12-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
3623-MSM12-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
3623-MSM12-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	INHA/9352
Immunogen	Recombinant human INHA protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	39.67kDa
Cellular Localization	Secreted
Species Reactivity	Human
Positive Control	Originally found in ovary (granulosa cells) and testis (Sertoli cells), but widely distributed in many tissues including brain and placenta

*Optimal dilution for a specific application should be determined.

Product Images for Inhibin, alpha (INHA) (Gonadal Cell Marker) Antibody

Specificity & Comments

Inhibins and activins inhibit and activate, respectively, the secretion of follitropin by the pituitary gland. Inhibins/activins are involved in regulating a number of diverse functions such as hypothalamic and pituitary hormone secretion, gonadal hormone secretion, germ cell development and maturation, erythroid differentiation, insulin secretion, nerve cell survival, embryonic axial development or bone growth, depending on their subunit composition. Inhibins appear to oppose the functions of activins., Inhibin A is a dimer of alpha/INHA and beta-A/INHBA that functions as a feedback regulator in the hypothalamic-pituitary-gonadal (HPG) axis. Inhibits the secretion of FSH from the anterior pituitary gland by acting on pituitary gonadotrope cells. Antagonizes activin A by binding to the proteoglycan, betaglycan, and forming a stable complex with and, thereby, sequestering type II activin receptors while excluding type I receptor., Inhibin B is a dimer of alpha and beta-B that plays a crucial role in the regulation of the reproductive system by inhibiting the secretion of follicle-stimulating hormone (FSH) from the anterior pituitary gland. Thereby, maintains reproductive homeostasis in both males and females. Acts as a more potent suppressor of FSH release than inhibin A (By similarity). Functions as competitive receptor antagonist binding activin type II receptors with high affinity in the presence of the TGF-beta type III coreceptor/TGFBR3L (PubMed:34910520).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.