

Hepatocyte growth factor receptor Antibody

Mouse Monoclonal Antibody [Clone MET/1871]

Catalog No	Format	Size
4233-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4233-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4233-MSM1-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	

Product Details	
Clone	MET/1871
Immunogen	Recombinant fragment of human c-MET protein; epitope is extracellular receptor of HGFR
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	155.54kDa
Cellular Localization	Membrane, Secreted
Species Reactivity	Human
Positive Control	Expressed in normal hepatocytes as well as in epithelial cells lining the stomach, the small and the large intestine

*Optimal dilution for a specific application should be determined.

Product Images for Hepatocyte growth factor receptor Antibody

Specificity & Comments

Receptor tyrosine kinase that transduces signals from the extracellular matrix into the cytoplasm by binding to hepatocyte growth factor/HGF ligand. Regulates many physiological processes including proliferation, scattering, morphogenesis and survival. Ligand binding at the cell surface induces autophosphorylation of MET on its intracellular domain that provides docking sites for downstream signaling molecules. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1, SRC, GRB2, STAT3 or the adapter GAB1. Recruitment of these downstream effectors by MET leads to the activation of several signaling cascades including the RAS-ERK, PI3 kinase-AKT, or PLCgamma-PKC. The RAS-ERK activation is associated with the morphogenetic effects while PI3K/AKT coordinates prosurvival effects. During embryonic development, MET signaling plays a role in gastrulation, development and migration of neuronal precursors, angiogenesis and kidney formation. During skeletal muscle development, it is crucial for the migration of muscle progenitor cells and for the proliferation of secondary myoblasts (By similarity). In adults, participates in wound healing as well as organ regeneration and tissue remodeling. Also promotes differentiation and proliferation of hematopoietic cells. May regulate cortical bone osteogenesis (By similarity)., (Microbial infection) Acts as a receptor for Listeria monocytogenes internalin InIB, mediating entry of the pathogen into cells.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.