



Interstitial collagenase Antibody

Mouse Monoclonal Antibody [Clone MMP1/4299]

Catalog No	Format	Size
4312-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4312-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4312-MSM3-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	MMP1/4299
Immunogen	Recombinant human MMP1 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	54.01kDa
Cellular Localization	Extracellular matrix, Extracellular space, Secreted
Species Reactivity	Human

**Optimal dilution for a specific application should be determined.*

Product Images for Interstitial collagenase Antibody

Specificity & Comments

Cleaves collagens of types I, II, and III at one site in the helical domain. Also cleaves collagens of types VII and X (PubMed:1645757, PubMed:2153297, PubMed:2557822). In case of HIV infection, interacts and cleaves the secreted viral Tat protein, leading to a decrease in neuronal Tat's mediated neurotoxicity (PubMed:16807369).

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.