ATP5F1A Antibody

Mouse Monoclonal Antibody [Clone ATP5A1/14170]

Catalog No	Format	Size
498-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
498-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
498-MSM4-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	ATP5A1/14170
Immunogen	Recombinant fragment (around aa 203-314) of human ATP5F1A protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	59.75kDa
Cellular Localization	Cell membrane, Mitochondrion, Mitochondrion inner membrane
Species Reactivity	Human
Positive Control	Fetal lung, heart, liver, gut and kidney

**Optimal dilution for a specific application should be determined.*

Product Images for ATP5F1A Antibody



Western blot analysis of skeletal muscle and stomach tissue lysates from human, mouse, rat, hamster, guinea pig and rabbit using ATP5A1 Mouse Monoclonal Antibody (ATP5A1/14170).



SDS-PAGE Analysis of Purified ATP5F1A Mouse Monoclonal Antibody (ATP5A1/14170). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

Subunit alpha, of the mitochondrial membrane ATP synthase complex (F(1)F(0) ATP synthase or Complex V) that produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain (Probable). ATP synthase complex consist of a soluble F(1) head domain - the catalytic core - and a membrane F(1) domain - the membrane proton channel (PubMed:37244256). These two domains are linked by a central stalk rotating inside the F(1) region and a stationary peripheral stalk (PubMed:37244256). During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation (Probable). In vivo, can only synthesize ATP although its ATP hydrolase activity can be activated artificially in vitro (By similarity). With the catalytic subunit beta (ATP5F1B), forms the catalytic core in the F(1) domain (PubMed:37244256). Subunit alpha does not bear the catalytic high-affinity ATP-binding sites (Probable). Binds the bacterial siderophore enterobactin and can promote mitochondrial accumulation of enterobactin-derived iron ions (PubMed:30146159).

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.