

## Phosphatidylcholine translocator ABCB4 Antibody

Mouse Monoclonal Antibody [Clone ABCB4/12631]

Catalog No	Format	Size
5244-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5244-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5244-MSM1-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details	
Clone	ABCB4/12631
Immunogen	Recombinant human ABCB4 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	141.52kDa
Cellular Localization	Apical cell membrane, Cell membrane, Clathrin-coated vesicle, Cytoplasm, Cytoplasmic vesicle, Membrane raft
Species Reactivity	Human

\*Optimal dilution for a specific application should be determined.

### Product Images for Phosphatidylcholine translocator ABCB4 Antibody

#### Specificity & Comments

Energy-dependent phospholipid efflux translocator that acts as a positive regulator of biliary lipid secretion. Functions as a floppase that translocates specifically phosphatidylcholine (PC) from the inner to the outer leaflet of the canalicular membrane bilayer into the canalculi of hepatocytes. Translocation of PC makes the biliary phospholipids available for extraction into the canalculi lumen by bile salt mixed micelles and therefore protects the biliary tree from the detergent activity of bile salts (PubMed:17523162, PubMed:21820390, PubMed:23468132, PubMed:24594635, PubMed:24723470, PubMed:24806754, PubMed:31873305, PubMed:7957936, PubMed:8898203, PubMed:9366571). Plays a role in the recruitment of phosphatidylcholine (PC), phosphatidylethanolamine (PE) and sphingomyelin (SM) molecules to nonraft membranes and to further enrichment of SM and cholesterol in raft membranes in hepatocytes (PubMed:23468132). Required for proper phospholipid bile formation (By similarity). Indirectly involved in cholesterol efflux activity from hepatocytes into the canalicular lumen in the presence of bile salts in an ATP-dependent manner (PubMed:24045840). Promotes biliary phospholipid secretion as canalculi-containing vesicles from the canalicular plasma membrane (PubMed:28012258, PubMed:9366571). In cooperation with ATP8B1, functions to protect hepatocytes from the deleterious detergent activity of bile salts (PubMed:21820390). Does not confer multidrug resistance (By similarity).

#### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

#### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

#### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.