

Peroxisome proliferator-activated receptor delta Antibody

Mouse Monoclonal Antibody [Clone PCR-P-PPARD-1D5]

Catalog No	Format	Size
5467-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5467-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5467-MSM4-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	

Product Details	
Clone	PCR-P-PPARD-1D5
Immunogen	Recombinant human PPARD protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2 / Kappa
Mol. Weight of Antigen	49.9kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	Ubiquitous with maximal levels in placenta and skeletal muscle

**Optimal dilution for a specific application should be determined.*

Product Images for Peroxisome proliferator-activated receptor delta Antibody

Specificity & Comments

Ligand-activated transcription factor key mediator of energy metabolism in adipose tissues (PubMed:35675826). Receptor that binds peroxisome proliferators such as hypolipidemic drugs and fatty acids. Has a preference for poly-unsaturated fatty acids, such as gamma-linoleic acid and eicosapentanoic acid. Once activated by a ligand, the receptor binds to promoter elements of target genes. Regulates the peroxisomal beta-oxidation pathway of fatty acids. Functions as transcription activator for the acyl-CoA oxidase gene. Decreases expression of NPC1L1 once activated by a ligand.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.