

BMI1 (Oncogene and Stem Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone PCR-P-BMI1-1G9]

Catalog No	Format	Size
648-MSM6-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
648-MSM6-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
648-MSM6-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	

Product Details

Clone	PCR-P-BMI1-1G9
Immunogen	Recombinant human BMI1 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2 / Kappa
Mol. Weight of Antigen	36.95kDa
Cellular Localization	Cytoplasm, Nucleus
Species Reactivity	Human

*Optimal dilution for a specific application should be determined.

Product Images for BMI1 (Oncogene and Stem Cell Marker) Antibody

Specificity & Comments

Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:15386022, PubMed:16359901, PubMed:16714294, PubMed:21772249, PubMed:25355358, PubMed:26151332, PubMed:27827373). The complex composed of RNF2, UB2D3 and BMI1 binds nucleosomes, and has activity only with nucleosomal histone H2A (PubMed:21772249, PubMed:25355358). In the PRC1-like complex, regulates the E3 ubiquitin-protein ligase activity of RNF2/RING2 (PubMed:15386022, PubMed:21772249, PubMed:26151332).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.