

## GLUT-1 (Tumor Progression and Mesothelioma Marker) Antibody

Mouse Monoclonal Antibody [Clone GLUT1/1973]

Catalog No	Format	Size
6513-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6513-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6513-MSM3-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details

Clone	GLUT1/1973
Immunogen	Recombinant fragment (around aa 203-305) of human SLC2A1 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b
Mol. Weight of Antigen	54.08kDa
Cellular Localization	Cell membrane, Melanosome, Photoreceptor inner segment
Species Reactivity	Human
Positive Control	Detected in erythrocytes (at protein level)

\*Optimal dilution for a specific application should be determined.

### Product Images for GLUT-1 (Tumor Progression and Mesothelioma Marker) Antibody

#### Specificity & Comments

Facilitative glucose transporter, which is responsible for constitutive or basal glucose uptake (PubMed:10227690, PubMed:10954735, PubMed:18245775, PubMed:19449892, PubMed:25982116, PubMed:27078104, PubMed:32860739). Has a very broad substrate specificity; can transport a wide range of aldoses including both pentoses and hexoses (PubMed:18245775, PubMed:19449892). Most important energy carrier of the brain: present at the blood-brain barrier and assures the energy-independent, facilitative transport of glucose into the brain (PubMed:10227690). In association with BSG and NXNL1, promotes retinal cone survival by increasing glucose uptake into photoreceptors (By similarity). Required for mesendoderm differentiation (By similarity).

#### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

#### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

#### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.