

Somatostatin (SST) Antibody

Mouse Monoclonal Antibody [Clone SST/16514]

Catalog No	Format	Size
6750-MSM7-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
6750-MSM7-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
6750-MSM7-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	SST/16514
Immunogen	Recombinant fragment (around aa 1-116) of human SST protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2c
Mol. Weight of Antigen	12.74kDa
Cellular Localization	Secreted
Species Reactivity	Human

*Optimal dilution for a specific application should be determined.

Product Images for Somatostatin (SST) Antibody

Specificity & Comments

Inhibits the secretion of pituitary hormones, including that of growth hormone/somatotropin (GH1), PRL, ACTH, luteinizing hormone (LH) and TSH. Also impairs ghrelin- and GnRH-stimulated secretion of GH1 and LH; the inhibition of ghrelin-stimulated secretion of GH1 can be further increased by neuronostatin. May enhance low-glucose-induced glucagon release by pancreatic alpha cells (By similarity). This effect may be mediated by binding to GPR107 and PKA activation (By similarity). May regulate cardiac contractile function (By similarity). May compromise cardiomyocyte viability (By similarity). In the central nervous system, may impair memory retention and may affect hippocampal excitability (By similarity). May also have anxiolytic and anorexigenic effects (By similarity). May play a role in arterial pressure regulation (By similarity). May inhibit basal, but not ghrelin- or GnRH-stimulated secretion of GH1 or LH, but does not affect the release of other pituitary hormones, including PRL, ACTH, FSH or TSH. Potentiates inhibitory action of somatostatin on ghrelin-stimulated secretion of GH1, but not that on GnRH-stimulated secretion of LH (PubMed:29615476).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.