



Wilm's Tumor 1 (WT1) (Wilm's Tumor & Mesothelial Marker) Antibody

Mouse Monoclonal Antibody [Clone WT1/7714]

Catalog No	Format	Size
7490-MSM14-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7490-MSM14-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7490-MSM14-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	WT1/7714
Immunogen	Recombinant human WT1 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	49.19kDa
Cellular Localization	Cytoplasm, Nucleolus, Nucleoplasm, Nucleus, Nucleus speckle
Species Reactivity	Human
Positive Control	Expressed in the kidney and a subset of hematopoietic cells

**Optimal dilution for a specific application should be determined.*

Product Images for Wilm's Tumor 1 (WT1) (Wilm's Tumor & Mesothelial Marker) Antibody

Specificity & Comments

Transcription factor that plays an important role in cellular development and cell survival (PubMed:7862533). Recognizes and binds to the DNA sequence 5'-GCG(T/G)GGGCG-3' (PubMed:17716689, PubMed:25258363, PubMed:7862533). Regulates the expression of numerous target genes, including EPO. Plays an essential role for development of the urogenital system. It has a tumor suppressor as well as an oncogenic role in tumor formation. Function may be isoform-specific: isoforms lacking the KTS motif may act as transcription factors (PubMed:15520190). Isoforms containing the KTS motif may bind mRNA and play a role in mRNA metabolism or splicing (PubMed:16934801). Isoform 1 has lower affinity for DNA, and can bind RNA (PubMed:19123921).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.