

Calcitonin receptor Antibody

Mouse Monoclonal Antibody [Clone CALCR/14449]

Catalog No	Format	Size
799-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
799-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
799-MSM2-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	CALCR/14449
Immunogen	Recombinant fragment (around aa 25-148) of human CALCR protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1
Mol. Weight of Antigen	55.34kDa
Cellular Localization	Cell membrane
Species Reactivity	Human

**Optimal dilution for a specific application should be determined.*

Product Images for Calcitonin receptor Antibody

Specificity & Comments

G protein-coupled receptor activated by ligand peptides amylin (IAPP), calcitonin (CT/CALCA) and calcitonin gene-related peptide type 1 (CGRP1/CALCA) (PubMed:35324283, PubMed:38603770). CALCR interacts with receptor-activity-modifying proteins RAMP1, 2 and 3 to form receptor complexes AMYR1, 2 and 3, respectively (PubMed:35324283, PubMed:38603770). IAPP, CT and CGRP1 activate CALCR and AMYRs with distinct modes of receptor activation resulting in specific phenotypes (PubMed:35324283, PubMed:38603770). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors. Activates cAMP-dependent pathway (PubMed:35324283, PubMed:7476993)., Non-functional protein. Unable to couple to G proteins and activate adenylyl cyclase (PubMed:7476993). Does not undergo receptor internalization following ligand binding (PubMed:7476993).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.