

Ligand-dependent corepressor Antibody

Mouse Monoclonal Antibody [Clone PCR-P-LCOR-1B1]

Catalog No	Format	Size
84458-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
84458-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
84458-MSM2-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	

Product Details

Clone	PCR-P-LCOR-1B1
Immunogen	Recombinant human LCOR protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2 / Kappa
Mol. Weight of Antigen	47.01kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	Ubiquitous

*Optimal dilution for a specific application should be determined.

Product Images for Ligand-dependent corepressor Antibody

Specificity & Comments

May act as transcription activator that binds DNA elements with the sequence 5'-CCCTATCGATCGATCTCTACCT-3' (By similarity). Repressor of ligand-dependent transcription activation by target nuclear receptors. Repressor of ligand-dependent transcription activation by ESR1, ESR2, NR3C1, PGR, RARA, RARB, RARG, RXRA and VDR.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.