

NFATC2-interacting protein Antibody

Mouse Monoclonal Antibody [Clone NFATC2IP/397]

Catalog No	Format	Size
84901-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
84901-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
84901-MSM1-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	

Product Details

Clone	NFATC2IP/397
Immunogen	Recombinant human NFATC2IP protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	45.82kDa
Cellular Localization	Cytoplasm, Nucleus
Species Reactivity	Human

*Optimal dilution for a specific application should be determined.

Product Images for NFATC2-interacting protein Antibody

Specificity & Comments

In T-helper 2 (Th2) cells, regulates the magnitude of NFAT-driven transcription of a specific subset of cytokine genes, including IL3, IL4, IL5 and IL13, but not IL2. Recruits PRMT1 to the IL4 promoter; this leads to enhancement of histone H4 'Arg-3'-methylation and facilitates subsequent histone acetylation at the IL4 locus, thus promotes robust cytokine expression (By similarity). Down-regulates formation of poly-SUMO chains by UBE2I/UBC9 (By similarity).

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.