

T-cell surface glycoprotein CD1c Antibody

Mouse Monoclonal Antibody [Clone CD1C/13378]

Catalog No	Format	Size
911-MSM13-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
911-MSM13-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
911-MSM13-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	CD1C/13378
Immunogen	Recombinant fragment (around aa 36-127) of human CD1C protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG
Mol. Weight of Antigen	37.65kDa
Cellular Localization	Cell membrane, Endosome membrane, Lysosome
Species Reactivity	Human
Positive Control	Expressed on cortical thymocytes, on certain T-cell leukemias, and in various other tissues

**Optimal dilution for a specific application should be determined.*

Product Images for T-cell surface glycoprotein CD1c Antibody

Specificity & Comments

Antigen-presenting protein that binds self and non-self lipid and glycolipid antigens and presents them to T-cell receptors on natural killer T-cells.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.