Recombinant CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody

Rabbit Monoclonal Antibody [Clone LFA2/8250R]

Catalog No	Format	Size
914-RBM19-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
914-RBM19-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
914-RBM19-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	LFA2/8250R
Immunogen	Recombinant human CD2 protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	39.45kDa
Cellular Localization	Cell membrane
Species Reactivity	Human
Positive Control	Expressed in natural killer cells (at protein level)

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody

Specificity & Comments

CD2 interacts with lymphocyte function-associated antigen CD58 (LFA-3) and CD48/BCM1 to mediate adhesion between T-cells and other cell types. CD2 is implicated in the triggering of T-cells, the cytoplasmic domain is implicated in the signaling function.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.