

Lymphocyte function-associated antigen 3 Antibody

Mouse Monoclonal Antibody [Clone CD58/1630]

Catalog No	Format	Size
965-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
965-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
965-MSM1-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	

Product Details

Clone	CD58/1630
Immunogen	Recombinant human CD58 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	28.15kDa
Cellular Localization	Cell membrane
Species Reactivity	Human

**Optimal dilution for a specific application should be determined.*

Product Images for Lymphocyte function-associated antigen 3 Antibody

Specificity & Comments

Ligand of the T-lymphocyte CD2 glycoprotein. This interaction is important in mediating thymocyte interactions with thymic epithelial cells, antigen-independent and -dependent interactions of T-lymphocytes with target cells and antigen-presenting cells and the T-lymphocyte rosetting with erythrocytes. In addition, the LFA-3/CD2 interaction may prime response by both the CD2+ and LFA-3+ cells.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.