

Ubiquitin-conjugating enzyme E2 R1 Antibody

Mouse Monoclonal Antibody [Clone CDC34/7505]

Catalog No	Format	Size
997-MSM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
997-MSM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
997-MSM5-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	CDC34/7505
Immunogen	Recombinant human CDC34 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2c / Kappa
Mol. Weight of Antigen	26.74kDa
Cellular Localization	Cytoplasm, Nucleus
Species Reactivity	Human
Positive Control	Expressed in testes during spermatogenesis to regulate repression of cAMP-induced transcription

*Optimal dilution for a specific application should be determined.

Product Images for Ubiquitin-conjugating enzyme E2 R1 Antibody



Western blot analysis of human brain and liver tissue lysates using CDC34 Mouse Monoclonal Antibody (CDC34/7505).



Western blot analysis of brain and liver tissue lysates from human, mouse, rat, hamster, guinea pig and rabbit using CDC34 Mouse Monoclonal Antibody (CDC34/7505).

Specificity & Comments

E2 ubiquitin-conjugating enzyme that accepts ubiquitin from an E1 ubiquitin-activating protein, and catalyzes its covalent attachment to other proteins by an E3 ubiquitin-protein ligase complex (PubMed:10329681, PubMed:17588522, PubMed:20061386, PubMed:38326650). In vitro catalyzes 'Lys-48'-linked polyubiquitination (PubMed:22496338). Cooperates with the E2 UBCH5C and the SCF(FBXW11) E3 ligase complex for the polyubiquitination of NFKBIA leading to its subsequent proteasomal degradation (PubMed:10329681, PubMed:10918611, PubMed:17698585). Performs ubiquitin chain elongation building ubiquitin chains from the UBE2D3-primed NFKBIA-linked ubiquitin. UBE2D3 acts as an initiator E2, priming the phosphorylated NFKBIA target at positions 'Lys-21' and/or 'Lys-22' with a monoubiquitin. Cooperates with the SCF(SKP2) E3 ligase complex to regulate cell proliferation through ubiquitination and degradation of MYBL2 and KIP1 (PubMed:10871850, PubMed:15652359, PubMed:19112177). Involved in ubiquitin conjugation and degradation of CREM isoform ICERIIgamma and ATF15 resulting in abrogation of ICERIIgamma- and ATF5-mediated repression of cAMP-induced transcription during both meiotic and mitotic cell cycles. Involved in the regulation of the cell cycle G2/M phase through its targeting of the WEE1 kinase for ubiquitination and degradation (PubMed:19126550). Also involved in the degradation of beta-catenin (PubMed:12037680). Is target of human herpes virus 1 protein ICP0, leading to ICP0-dependent dynamic interaction with proteasomes (PubMed:11805320, PubMed:12060736).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.